

**Modified T-stenting technique with crushing for bifurcation lesions: immediate results and 30-day outcome.** Colombo, A., G. Stankovic, et al. Catheter Cardiovasc Interv (2003).**60**(2): 145-51  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=14517916](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14517916)

**Dominant contribution of negative remodeling to development of significant coronary bifurcation narrowing.** Fujii, K., Y. Kobayashi, et al. Am J Cardiol (2003).**92**(1): 59-61  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12842248](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12842248)

**A new dedicated stent and delivery system for the treatment of bifurcation lesions: preliminary experience.** Toutouzas, K., G. Stankovic, et al. Catheter Cardiovasc Interv (2003).**58**(1): 34-42  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12508194](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12508194)