

Effectiveness of elective stenting of common carotid artery lesions in preventing stroke. Chio, F. L., M. W. Liu, et al. Am J Cardiol (2003).**92**(9): 1135-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14583376

Peripheral vascular disease and one-year mortality following percutaneous coronary revascularization. Chiu, J. H., E. J. Topol, et al. Am J Cardiol (2003).**92**(5): 582-3

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12943879

Randomized study to compare balloon angioplasty and elective stent implantation in venous bypass grafts: the Venestent study. Hanekamp, C. E., J. J. Koolen, et al. Catheter Cardiovasc Interv (2003).**60**(4): 452-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14624420

Predictors of restenosis after successful carotid artery stenting. Khan, M. A., M. W. Liu, et al. Am J Cardiol (2003).**92**(7): 895-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14516904

Carotid sinus reactions during carotid artery stenting: predictors, incidence, and influence on clinical outcome. Leisch, F., K. Kerschner, et al. Catheter Cardiovasc Interv (2003).**58**(4): 516-23

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12652504

Statin use and leg functioning in patients with and without lower-extremity peripheral arterial disease. McDermott, M. M., J. M. Guralnik, et al. Circulation (2003).**107**(5): 757-61

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12578881

Cholesterol reduction with atorvastatin improves walking distance in patients with peripheral arterial disease. Mohler, E. R., 3rd, W. R. Hiatt, et al. Circulation (2003).**108**(12): 1481-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12952839

Endovascular intervention of aortoiliac occlusive disease in high-risk patients using the kissing stents technique: Long-term results. Mouanoutoua, M., R. Maddikunta, et al. Catheter Cardiovasc Interv (2003).**60**(3): 320-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14571480

Regional angiogenesis with vascular endothelial growth factor in peripheral arterial disease: a phase II randomized, double-blind, controlled study of adenoviral delivery of vascular endothelial growth factor 121 in patients with disabling intermittent claudication. Rajagopalan, S., E. R. Mohler, 3rd, et al. Circulation (2003).**108**(16): 1933-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14504183

Focal ischemia of the brain after neuroprotected carotid artery stenting.

Schluter, M., T. Tubler, et al. *J Am Coll Cardiol* (2003).**42**(6): 1007-13

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=13678921

Endovascular stent implantation in patients with stenotic

aortoarteriopathies: early and medium-term results. Siwik, E. S., S. B. Perry, et al. *Catheter Cardiovasc Interv* (2003).**59**(3): 380-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12822165

Self- and balloon-expandable stent implantation for severe native coarctation of aorta in adults. Tyagi, S., S. Singh, et al. *Am Heart J* (2003).**146**(5): 920-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14597945

The adult patient with native coarctation of the aorta: balloon angioplasty or primary stenting? Zabal, C., F. Attie, et al. *Heart* (2003).**89**(1): 77-83

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12482798

Predictors of improved renal function after percutaneous stent-supported angioplasty of severe atherosclerotic ostial renal artery stenosis. Zeller, T., U. Frank, et al. *Circulation* (2003).**108**(18): 2244-9

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14557357

Predictors of Improved Renal Function After Percutaneous Stent-Supported Angioplasty of Severe Atherosclerotic Ostial Renal Artery Stenosis. Zeller, T., U. Frank, et al. *Circulation* (2003).**108**(18): 2244-2249

<http://circ.ahajournals.org/cgi/content/abstract/108/18/2244>

Cutting balloon angioplasty of the popliteal and infrapopliteal vessels for symptomatic limb ischemia. Ansel, G. M., N. S. Sample, et al. *Catheter Cardiovasc Interv* (2004).**61**(1): 1-4

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14696150

Subclavian artery stenting: factors influencing long-term outcome. Bates, M. C., M. Broce, et al. *Catheter Cardiovasc Interv* (2004).**61**(1): 5-11

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14696151